

Consultant Level Report for:

Mr Siddharth Prashant Govilkar GMC Number: 7016356 NJR Number: 7528

For the Period to 31 March 2023

The BOA and the NJR are agreed that this document should be used to inform each surgeon's annual appraisal. The document should be shown to, and discussed with, the appraiser or with another arthroplasty surgeon if the appraiser is of another speciality. This is in keeping with the policy of the GMC which states that outcomes data from any national audit should be utilised for the purpose of annual appraisal.

Contents

Organisation Summary
Data Quality
Hips
Knees
Appendix 1: SRR Funnel Plots (as Consultant in Charge) - Hips
Appendix 1: SRR Funnel Plots (as Consultant in Charge) - Knees
Appendix 2: SRR Funnel Plots (as Lead Surgeon) - Hips
Appendix 2: SRR Funnel Plots (as Lead Surgeon) - Knees
Appendix 3: 90 day SMR (as Consultant in Charge) - Hips
Appendix 3: 90 day SMR (as Consultant in Charge) - Knees
Appendix 4: 90 day SMR (as Lead Surgeon) - Hips
Appendix 4: 90 day SMR (as Lead Surgeon) - Knees

This report has been produced by the National Joint Registry (NJR). It represents all activity recorded in the NJR, in the name of the selected surgeon (as Consultant in Charge), up to the specified period.

This report is made available to the named surgeon for personal review, to share with colleagues, and to be used in consultant revalidation. The named surgeon in this report is free to share this report as they choose subject to data protection legislation and local governance permissions. Please note that some tables contain record level data and the identities of third party surgeons and so care should be taken when sharing or publishing.

Constraints

This report reflects data reported in the NJR. Missing data and issues with the quality of data recorded within the NJR may impact the results shown and may make the results of either a unit or surgeon seem better OR worse than should be the case. You should consider the following in assessing the data quality of the report:

• Consent – an assessment of the proportion of patients at your trust who provide consent for their details to be recorded within the NJR. Without consent, it is not possible to link primary and revision procedures in the calculation of revision rates.

• NJR Compliance – The percentage of all total joint procedures for your trust that have been entered into the NJR within any given period compared with the number of procedures submitted to HES and/or PEDW.

Further Information

Further analysis of this data is possible through NJR Connect – Data Services, https://platform.njrcentre.org.uk/

For further information please visit the NJR website at <u>www.njrcentre.org.uk</u> or contact the NJR Service Desk on 0845 345 9991 or <u>enquiries@njrcentre.org.uk</u>.

Organisation Summary

In this Section : Total primary and revision activity recorded for the surgeon (as Consultant in Charge) on the NJR over a 12 and 36 month period, the joint type, and the hospital in which the operation was performed.

12 month activity: for the period 1 Apr 2022 - 31 Mar 2023

Organisation Type	Unit	Hip	Knee	Ankle	Elbow	Shoulder	Total	% of activity
Independent Sector	Nuffield Health North Staffordshire Hospital	36	32	0	0	0	68	52%
NHS	County Hospital	33	12	0	0	0	45	35%
NHS	Royal Stoke University Hospital	17	0	0	0	0	17	13%
Total		86	44	0	0	0	130	

36 month activity: for the period 1 Apr 2020 - 31 Mar 2023

Organisation Type	Unit	Hip	Knee	Ankle	Elbow	Shoulder	Total	% of activity
Independent Sector	Nuffield Health North Staffordshire Hospital	36	32	0	0	0	68	52%
NHS	County Hospital	33	12	0	0	0	45	35%
NHS	Royal Stoke University Hospital	17	0	0	0	0	17	13%
Total		86	44	0	0	0	130	



Data Quality

In this Section : Measure of consent is the proportion of all patients who agree to have their data stored in the NJR. Where consent is less than average, this may mean that not all data for the surgeon is loaded on the NJR, or that missing data for the surgeon reduces the reliability of indicators and outcome measures. Counts are for the period 1 Apr 2022 – 31 Mar 2023.

NJR Consent:

Unit level measures where this surgeon has activity:


Measure	NJR Consent	Organisation Ty	pe Unit	NJR Consent
National Average Rate	93.51%	Independent Sect	tor Nuffield Health North Staffordshire Hospital	100.00%
This surgeon's rate	100.00%	NHS	County Hospital	100.00%
		NHS	Royal Stoke University Hospital	100.00%



Hips

Hips – Recorded Activity

In this Section : Volume, procedure type and hip articulation undertaken by the surgeon over a 36 month period, showing year on year trend. Note that Debridement and Implant Retention (DAIR) procedures that did not involve a modular exchange are not included in the count of revision procedures and are shown separately.



Procedure Type	Apr22- Mar23
Primary Cemented	80
Primary Cementless	1
Revision	5
Total	86

Hip ArticulationApr22-
Mar23Ceramic On Poly63Metal On Poly19Total82

DAIR without Modular Exchange

None Recorded

Dual Consultant Operating Cases


This table shows the grade of the first assistant surgeon on all procedures in which the consultant is named as having participated in the latest 12 month and 36 month periods.

	First Assistant Grade						
Consultant in Charge/Lead Surgeon Combination	Cons	ultant	Other				
	Last Year	Last 3 Years	Last Year	Last 3 Years			
Mr Siddharth Prashant Govilkar both CiC and Lead Surgeon	3	3	83	83			
Mr Siddharth Prashant Govilkar CiC with another consultant as Lead Surgeon	0	0	0	0			
Mr Siddharth Prashant Govilkar CiC/with a Non-Consultant Grade as Lead Surgeon	0	0	0	0			
Consultant Grade CiC with Mr Siddharth Prashant Govilkar as Lead Surgeon	22	26	0	0			



Hips – Patient Profile

In this Section : The profile of hip patients operated on in the name of the surgeon (as Consultant in Charge) over the 12 month period 1 Apr 2022 – 31 Mar 2023.



BMI						
Patient Median BMI						
National	28.00					
Surgeon	29.00					

	Patient Mean Age
National	68.90
Surgeon	68.10



Hips

Outcomes following Primary Hip Surgery

In this Section : Quality and outcome measures for patients receiving primary hip replacement surgery by this surgeon based on an analysis of the most recent ten years' of data. This is a change to previous reports where the analysis has been based on all data held by the NJR, i.e. 'life of registry'. The SRR and SMR data are also available as funnel plots: see Appendices 1, 2, 3 and 4.

Indicator Set	Indicator	Linkable Primaries	Expected Events	Observed Events	Outside 99.8% control limts A This Surgeon 1 2 3	Inside 99.8% control limts 4 5
Mortality	Hip All – Last Five Years	36	0.05	0		
Revision	Hip All - Last Ten Years	43	0.04	0	4	
Revision	Hip All - Last Five Years	43	0.04	0		
Revision	Cemented Hip Procedures - Last Ten Years	43	0.03	0	4	
Revision	Cementless Hip Procedures - Last Ten Years	N/A	N/A	N/A	None Record	∋d
Revision	Hip Resurfacing - Last Ten Years	N/A	N/A	N/A	None Record	ed
Revision	Hybrid Hip Procedures - Last Ten Years	N/A	N/A	N/A	None Record	ed

Definitions

Mortality : Patient death (for whatever reason) within 90 days of the procedure having taken place.

The expected number of events in the table above are based on the number of primary procedures performed, and have been calculated from national average figures. The expected number has been adjusted to take into account selected patient variables such as age, gender, and ASA grade. The charts to the right depict the statistical significance of any difference between the observed number of events and corresponding expected number. There is a less than 0.1% chance that any individual surgeon will fall in the "better than expected" zone by chance alone, and similarly less than 0.1% chance of falling in the "worse than expected" zone. Note that depiction as an outlier does not constitute proof of over or under performance; some variation could be attributable to patient related (or other) risk factors that are not included in the adjustment model.

When calculating the 90 Day SMR, some primary procedures are excluded. For hips, this is procedures with indications of trauma or metastatic cancer/malignancy. For knees, it is procedures with trauma as an indication for implantation.

Note : Total number of procedures shown for Hip All indicators will include any Metal-on-Metal undertaken, but these procedures are not shown as a separate sub-category. The total number of procedures shown may, therefore, not be equal to the sum of the listed sub-categories.



National Joint Registry www.njrcentre.org.uk Working for patients, driving forward quality

6

Hip – Revision

In this Section: Based on the most recent ten years of data, the tables below may contain cases excluded from outcome analysis presented in the charts e.g. failed hemiarthroplasty cases. Tables containing record level data are included in Appendix 5.

Count of Revised Primaries by Year

This table shows, by year, the number of primaries recorded and the number of those primaries which have subsequently been revised.

None recorded

Unadjusted Revision Rate (all NJR)

Revision Period / Revision in	No of Recorded Primaries	No of Attributable Revisions	Unadjusted Revision Rate	National Average
1 Year	0	0	0.00%	0.73%
3 Years	0	0	0.00%	1.31%
5 Years	0	0	0.00%	1.92%



Hips – Implant Usage

The way in which ODEP applies ratings has recently changed, such that ratings now apply to 'constructs' rather than individual devices. The NJR is currently undertaking work to populate the new information so that it can be used for all future ODEP reporting. As a result, the report 'Implant Usage by ODEP Rating' will not be available in this format until next year.

Implant brand usage in primary hip replacement

The table compares, for the latest five years, the number of brands used in primary hip replacement procedures compared to the average for all surgeons in the NJR over the same period. The number of procedures for the latest five years is also included.


The brands are as listed in the NJR Annual Report. For further information, please access the Implant Library available on NJR Connect - Data Services, https://platform.njrcentre.org.uk/.

Number of procedures	Average per surgeon (all NJR)	Number of brands used	Average brand usage per surgeon (all NJR)
81	187	9	7



Knees – Recorded Activity

In this Section : Volume, procedure type undertaken by the surgeon over a 36 month period, showing year on year trend. Note that Debridement and Implant Retention (DAIR) procedures that did not involve a modular exchange are not included in the count of revision procedures and are shown separately.



Procedure Type	Apr22-Mar23
Primary Total Knee Replacement	44
Total	44

Knee Constraint	Apr22- Mar23
Posterior Stabilised Fixed	1
Unconstrained Fixed	43
Total	44

DAIR without Modular Exchange

None Recorded

Unicondylar Knee Replacement

The following table shows the number of unicondylar knee replacements undertaken in the previous two financial years. However, where both 'lateral' and 'medial' are indicated for a 'Multi-compartmental Knee Replacement', this is included as a single UKR.

None Recorded



Dual Consultant Operating Cases


This table shows the grade of the first assistant surgeon on all procedures in which the consultant is named as having participated in the latest 12 month and 36 month periods.

	First Assistant Grade					
Consultant in Charge/Lead Surgeon Combination	Cons	sultant	Other			
	Last Year	Last 3 Years	Last Year	Last 3 Years		
Mr Siddharth Prashant Govilkar both CiC and Lead Surgeon	1	1	43	43		
Mr Siddharth Prashant Govilkar CiC with another consultant as Lead Surgeon	0	0	0	0		
Mr Siddharth Prashant Govilkar CiC/with a Non-Consultant Grade as Lead Surgeon	0	0	0	0		
Consultant Grade CiC with Mr Siddharth Prashant Govilkar as Lead Surgeon	46	76	2	3		



Knees - Patient Profile

In this Section : The profile of knee patients operated on in the name of the surgeon (as Consultant in Charge) over the 12 month period 1 Apr 2022 – 31 Mar 2023.



BMI			
	Patient Median BMI		
National	30.00		
Surgeon	31.00		

	Patient Mean Age		
National	69.37		
Surgeon	71.61		



Outcomes following Primary Knee Surgery

In this Section : Quality and outcome measures for patients receiving primary knee replacement surgery by this surgeon based on an analysis of the most recent ten years' of data. This is a change to previous reports where the analysis has been based on all data held by the NJR, i.e. 'life of registry'. The SRR and SMR data are also available as funnel plots: see Appendices 1, 2, 3 and 4.

Indicator Set Mortality	Indicator Knee All – Last Five Years	Linkable Primaries 27	Expected Events	Observed Events	0	Outside 99.8% ontrol limts 1	•	This Su 2	irgeon 3	Insie 99.8 cont lim 4	8% rol
Revision	Knee All - Last Ten Years	27	0.02	0		4					
Revision	Total Knee Replacement - Last Ten Years	27	0.02	0							
Revision	Unicondylar Knee Procedures - Last Ten Years	N/A	N/A	N/A			Νοι	ne Ree	corded		
Revision	Patello-Femoral Knee Procedures - Last Ten Years	N/A	N/A	N/A			Noi	ne Ree	corded		
Revision	Knee All - Last Five Years	27	0.02	0		4					
Revision	Total Knee Replacement - Last Five Years	27	0.01	0							
Revision	Unicondylar Knee Procedures - Last Five Years	N/A	N/A	N/A			Noi	ne Ree	corded		

Definitions

Mortality : Patient death (for whatever reason) within 90 days of the procedure having taken place.

The expected number of events in the table above are based on the number of primary procedures performed, and have been calculated from national average figures. The expected number has been adjusted to take into account selected patient variables such as age, gender, and ASA grade. The charts to the right depict the statistical significance of any difference between the observed number of events and corresponding expected number. There is a less than 0.1% chance that any individual surgeon will fall in the "better than expected" zone by chance alone, and similarly less than 0.1% chance of falling in the "worse than expected" zone. Note that depiction as an outlier does not constitute proof of over or under performance; some variation could be attributable to patient related (or other) risk factors that are not included in the adjustment model.

When calculating the 90 Day SMR, some primary procedures are excluded. For hips, this is procedures with indications of trauma or metastatic cancer/malignancy. For knees, it is procedures with trauma as an indication for implantation.



Knee – Revision

In this Section: Based on the most recent ten years' of data. Tables containing record level data are included in Appendix 5.

Count of Revised Primaries by Year

This table shows, by year, the number of primaries recorded and the number of those primaries which have subsequently been revised.

None recorded

Unadjusted Revision Rate (all NJR)

Revision Period / Revision in	No of Recorded Primaries	No of Attributable Revisions	Unadjusted Revision Rate	National Average
1 Year	0	0	0.00%	0.46%
3 Years	0	0	0.00%	1.53%
5 Years	0	0	0.00%	2.32%



Knees – Implant Usage

The way in which ODEP applies ratings has recently changed, such that ratings now apply to 'constructs' rather than individual devices. The NJR is currently undertaking work to populate the new information so that it can be used for all future ODEP reporting. As a result, the report 'Implant Usage by ODEP Rating' will not be available in this format until next year.

Implant brand usage in primary knee replacement

The table compares, for the latest five years, the number of brands used in primary knee replacement procedures compared to the average for all surgeons in the NJR over the same period. The number of procedures for the latest five years is also included.


The brands are as listed in the NJR Annual Report. For further information, please access the Implant Library available on NJR Connect - Data Services, https://platform.njrcentre.org.uk/.


Number of procedures	Average per surgeon (all NJR)	Number of brands used	Average brand usage per surgeon (all NJR)
44	222	2	4




Hips

Appendix 1 : SRR Funnel Plots (as Consultant in Charge) - Hips









Hips


Appendix 1 : SRR Funnel Plots (as Consultant in Charge) - Hips






Appendix 1 : SRR Funnel Plots (as Consultant in Charge) - Knees








Appendix 1 : SRR Funnel Plots (as Consultant in Charge) - Knees

Standardised Revision Ratio Funnel Plot Representation: This section illustrates your Standardised Revision Ratio in the form of a funnel plot, based on the most recent ten years' of data. This is an alternative method of displaying the values shown within the main report for this indicator, and includes plots for all other surgeons.



Unicondylar Knee procedures - Last Ten years

None Recorded


Patello-femoral Knee procedures - Last Ten years

None Recorded

Data — Upper 99.8% — Upper 95% — Lower 95% — Lower 99.8%

patients, driving

Total Knee replacement - Last Five Years






Appendix 1 : SRR Funnel Plots (as Consultant in Charge) - Knees

Standardised Revision Ratio Funnel Plot Representation: This section illustrates your Standardised Revision Ratio in the form of a funnel plot, based on the most recent ten years' of data. This is an alternative method of displaying the values shown within the main report for this indicator, and includes plots for all other surgeons.

Unicondylar Knee procedures - Last Five Years


None Recorded








Hips

Appendix 2 : SRR Funnel Plots (as Lead Surgeon) - Hips










Hips

Appendix 2 : SRR Funnel Plots (as Lead Surgeon) - Hips


Standardised Revision Ratio Funnel Plot Representation: This section illustrates your Standardised Revision Ratio in the form of a funnel plot, based on the most recent ten years' of data. This is an alternative method of displaying the values shown within the main report for this indicator, and includes plots for all other surgeons.

5 4 Standardised Revision Ratio 3 2 1 0 5 0 10 15 20 Number of Expected Revisions

onal Joint Registry



Hybrid Hip procedures (from 2 linkable primary procedures)



Cemented Hip procedures (from 64 linkable primary procedures) Cementless Hip procedures (from 3 linkable primary procedures)




Data


Nati


www.njrcentre.org.uk


Appendix 2 : SRR Funnel Plots (as Lead Surgeon) - Knees

Standardised Revision Ratio Funnel Plot Representation: This section illustrates your Standardised Revision Ratio in the form of a funnel plot, based on the most recent ten years' of data. This is an alternative method of displaying the values shown within the main report for this indicator, and includes plots for all other surgeons.










Knee all - Last Five years (from 97 linkable primary procedures)


National Joint Registry www.njrcentre.org.uk Working for patients, driving forward quality

Appendix 2 : SRR Funnel Plots (as Lead Surgeon) - Knees

Standardised Revision Ratio Funnel Plot Representation: This section illustrates your Standardised Revision Ratio in the form of a funnel plot, based on the most recent ten years' of data. This is an alternative method of displaying the values shown within the main report for this indicator, and includes plots for all other surgeons.



Unicondylar Knee procedures - Last Ten years (from 1 linkable primary procedures)




Patello-femoral Knee procedures - Last Ten years

None Recorded

Upper 99.8% ____ Upper 95% ____ Lower 95% ____ Lower 99.8%

atients, driving

Total Knee replacement - Last Five Years (from 96 linkable primary procedures)







Data

Appendix 2 : SRR Funnel Plots (as Lead Surgeon) - Knees

Standardised Revision Ratio Funnel Plot Representation: This section illustrates your Standardised Revision Ratio in the form of a funnel plot, based on the most recent ten years' of data. This is an alternative method of displaying the values shown within the main report for this indicator, and includes plots for all other surgeons.

Unicondylar Knee procedures - Last Five Years (from 1 linkable primary procedures)









VJR

Hips

Appendix 3 : 90 day SMR (as Consultant in Charge) - Hips

Standardised Mortality Ratio Funnel Plot Representation: This section illustrates your Standardised Mortality Ratio (SMR) in the form of a funnel plot, based on the most recent five years' of data. It shows the SMR for mortality events occurring within 90 days of the primary procedure. This is an alternative method of displaying the values shown within the main report for this indicator and includes plots for all other surgeons.





Data — Upper 99.8% — Upper 95% — Lower 95% — Lower 99.8% • Outlier •



Mar

2023

Mar

2022

•

Mar

2021

•

Mar

2020


•


Mar

2019

Appendix 3 : 90 day SMR (as Consultant in Charge) - Knees

Standardised Mortality Ratio Funnel Plot Representation: This section illustrates your Standardised Mortality Ratio (SMR) in the form of a funnel plot, based on the most recent five years' of data. It shows the SMR for mortality events occurring within 90 days of the primary procedure. This is an alternative method of displaying the values shown within the main report for this indicator and includes plots for all other surgeons.





Data — Upper 99.8% — Upper 95% — Lower 95% — Lower 99.8% • Outlier • Mar 2023 • Mar 2021



Mar

2020

•


Mar

2019

Hips


Appendix 4 : 90 day SMR (as Lead Surgeon) - Hips

Standardised Mortality Ratio Funnel Plot Representation: This section illustrates your Standardised Mortality Ratio (SMR) in the form of a funnel plot, based on the most recent five years' of data. It shows the SMR for mortality events occurring within 90 days of the primary procedure. This is an alternative method of displaying the values shown within the main report for this indicator and includes plots for all other surgeons.



www.njrcentre.org.uk

Working for patients, driving forward quality





Appendix 4 : 90 day SMR (as Lead Surgeon) - Knees

Standardised Mortality Ratio Funnel Plot Representation: This section illustrates your Standardised Mortality Ratio (SMR) in the form of a funnel plot, based on the most recent five years' of data. It shows the SMR for mortality events occurring within 90 days of the primary procedure. This is an alternative method of displaying the values shown within the main report for this indicator and includes plots for all other surgeons.

